Journal of MOLECULAR BIOLOGY

EDITOR-IN-CHIEF
J. C. KENDREW

ASSISTANT EDITOR
S. BRENNER

EDITORIAL BOARD
C. R. Cantor
P. Chambon
W. W. Franke
M. Gellert
G. A. Gilbert
M. Gottesman
I. Herskowitz
R. Huber
H. E. Huxley
A. Klug
V. Luzzati
J. H. Miller
M. F. Moody
K. Simons

VOLUME 173

15 February to 15 March 1984

3

ACADEMIC PRESS
London New York San Francisco

Copyright © 1984, by Academic Press Inc. (London) Ltd.

ALL RIGHTS RESERVED

No part of this volume may be reproduced in any form, by photostat, microfilm, or any other means, without written permission from the publishers.

Printed in Great Britain

CONTENTS OF VOLUME 173

NUMBER 1, 15 FEBRUARY 1984

FAGRELIUS, T. J. and LIVINGSTON, D. M., Location of DNAase I Sensitive Cleavage Sites in the Yeast 2 μm Plasmid DNA Chromosome . . 1

MATSUBARA, I., GOLDMAN, Y. E. and SIMMONS, R. M., Changes in the Lateral Filament Spacing of Skinned Muscle Fibres when Cross-bridges Attach 15

Appendix 30

UENO, H. and HARRINGTON, W. F., An Enzyme-probe Method to Detect Structural Changes in the Myosin Rod 35

BOBST, A. M., KAO, S.-C., TOPPIN, R. C., IRELAND, J. C. and THOMAS, I. E., Dipsticking the Major Groove of DNA with Enzymatically Incorporated Spin-labeled Deoxyuridines by Electron Spin Resonance Spectroscopy 63

HOROWITZ, D. S. and WANG, J. C., Torsional Rigidity of DNA and Length Dependence of the Free Energy of DNA Supercoiling 75

SIGRIST, H., ALLEGRINI, P. R., STAUFFER, K., SCHALLER, J., ABDULAEV, N. G., RICKLI, E. E. and ZAHLER, P., Group-directed Modification of Bacteriorhodopsin by Arylisothiocyanates. Labeling, Identification of the Binding Site and Topology 93

GIOMETTI, C. S. and ANDERSON, N. L., An Electrophoretic Variant of a Human Fibroblast Protein with Characteristics of Smooth Muscle Tropomyosin 109

LETTERS TO THE EDITOR

KOYASU, S. and SHIRAKIHARA, Y., *Caulobacter crescentus* Flagellar Filament has a Right-handed Helical Form 125

LÜNSDORF, H., EHRIG, K., FRIEDL, P. and SCHAIRER, H. U., Use of Monoclonal Antibodies in Immuno-electron Microscopy for the Determination of Subunit Stoichiometry in Oligomeric Enzymes. There are Three α-Subunits in the F_1-ATPase of *Escherichia coli* . 131

POGLITSCH, A., KREMER, F., GENZEL, L., Picosecond Relaxations in Hydrated Lysozyme Observed by mm-Wave Spectroscopy . . 137

NUMBER 2, 25 FEBRUARY 1984

PAGÈS, M. J. M. and ROIZÈS, G. P., Nature and Organization of the Sequence Variations in the Long-range Periodicity Calf Satellite DNA I 143

ALEXANDER, F., YOUNG, P. R. and TILGHMAN, S. M., Evolution of the Albumin:α-Fetoprotein Ancestral Gene from the Amplification of a 27 Nucleotide Sequence 159

Appendix: YOUNG, P. R. and LITWIN, S., Determination of the Likelihood of Repetitive Sequences in the α-Fetoprotein Gene . 174

WU, T.-H., WOOD, D. L., STEIN, P. L. and COMER, M. M., Transcription of a Gene Cluster Coding for Two Aminoacyl-tRNA Synthetases and an Initiation Factor in *Escherichia coli* 177

FULFORD, W. and MODEL, P., Specificity of Translational Regulation by Two DNA-binding Proteins 211

RICHTER, J. D., SMITH, L. D., ANDERSON, D. M. and DAVIDSON, E. H., Interspersed Poly(A) RNAs of Amphibian Oocytes are not Translatable 227

VON HEIJNE, G., How Signal Sequences Maintain Cleavage Specificity. . 243

TAKEYA, M., KLIER, F. G. and FISHMAN, W. H., Molecular Dimensions of the SS and FF Phenotypes of Human Placental Alkaline Phosphatase. Rotary Shadowing and Negative Staining Electron Microscope Measurements 253

LETTER TO THE EDITOR

MOFFATT, B. A., DUNN, J. J. and STUDIER, F. W., Nucleotide Sequence of the Gene for Bacteriophage T7 RNA Polymerase . . . 265

ERRATUM 271

NUMBER 3, 5 MARCH 1984

WOOD, R. D., SKOPEK, T. R. and HUTCHINSON, F., Changes in DNA Base Sequence Induced by Targeted Mutagenesis of Lambda Phage by Ultraviolet Light 273

WOOD, R. D. and HUTCHINSON, F., Non-targeted Mutagenesis of Unirradiated Lambda Phage in *Escherichia coli* Host Cells Irradiated with Ultraviolet Light 293

ABELES, A. L., SNYDER, K. M. and CHATTORAJ, D. K., P1 Plasmid Replication: Replicon Structure 307

RICHARDS, O. C., MARTIN, S. C., JENSE, H. G. and EHRENFELD, E., Structure of Poliovirus Replicative Intermediate RNA. Electron Microscope Analysis of RNA Cross-linked *in Vivo* with Psoralen Derivative 325

WILLIAMSON, M. P., MARION, D. and WÜTHRICH, K., Secondary Structure in the Solution Conformation of the Proteinase Inhibitor IIA from Bull Seminal Plasma by Nuclear Magnetic Resonance 341

HOLBROOK, S. R. and KIM, S.-H., Local Mobility of Nucleic Acids as Determined from Crystallographic Data. I. RNA and *B* Form DNA . 361

LETTER TO THE EDITOR

AVOLIO, J., LEBDUSKA, S. and SATIR, P., Dynein Arm Substructure and the Orientation of Arm–Microtubule Attachments 389

NUMBER 4, 15 MARCH 1984

BISHOP, C., GUELLAEN, G., GELDWERTH, D., FELLOUS, M. and WEISSENBACH, J., Extensive Sequence Homologies Between Y and Other Human Chromosomes 403

HELLER, D., JACKSON, M. and LEINWAND, L., Organization and Expression of Non-Alu Family Interspersed Repetitive DNA Sequences in the Mouse Genome 419

RALEIGH, E. A. and KLECKNER, N., Multiple IS*10* Rearrangements in *Escherichia coli* 437

Appendix 453

KATO, S., OKAMOTO, M. and ASAKURA, S., Polymorphic Transition of the Flagellar Polyhook from *Escherichia coli* and *Salmonella typhimurium* . 463

MONTEILHET, C., BLOW, D. M. and BRICK, P. Interaction of Crystalline Tyrosyl-tRNA Synthetase with Adenosine, Adenosine Monophosphate, Adenosine Triphosphate and Pyrophosphate in the Presence of Tyrosinol 477

TAYLOR, W. R. and THORNTON, J. M., Recognition of Super-secondary Structure in Proteins 487

Appendix: TAYLOR, W. R., An Algorithm to Compare Secondary Structure Predictions 512

LETTERS TO THE EDITOR

RASHIN, A. A. and HONIG, B., On the Environment of Ionizable Groups in Globular Proteins 515

AHMED, A., A Deletion Analysis of Transposon Tn*9* 523

MÉRIC, C., DARLIX, J.-L. and SPAHR, P.-F., It is Rous Sarcoma Virus Protein P12 and not P19 that Binds Tightly to Rous Sarcoma Virus RNA. 531

Beck, E., Jorcano, J. L. and Alonso, A., *Drosophila melanogaster* U1 and U2 Small Nuclear RNA Genes Contain Common Flanking Sequences 539

Erratum 543

Index to Volume 173

Abdulaev, N. G. (see also Sigrist, H.), 93
Abeles, A. L., Snyder, K. M. & Chattoraj, D. K., P1 plasmid replication: replicon structure, 307
Adenosine, adenosine monophosphate, adenosine triphosphate and pyrophosphate, interaction with crystalline tyrosyl-tRNA synthetase in the presence of tyrosinol, 477
Ahmed, A., A deletion analysis of transposon Tn*9*, 523
Albumin: α-fetoprotein ancestral gene, evolution from the amplification of a 27 nucleotide sequence, 159
Alexander, F., Young, P. R. & Tilghman, S. M., Evolution of the albumin : α-fetoprotein ancestral gene from the amplification of a 27 nucleotide sequence, 159
Algorithm to compare secondary structure predictions, 512
Allegrini, P. R. (see also Sigrist, H.), 93
Alonso, A. (see also Beck, E.), 539
Aminoacyl-tRNA synthetases, two, and an initiation factor in *Escherichia coli*, transcription of the coding gene cluster, 177
AMP, interaction with crystalline tyrosyl-tRNA synthetase in the presence of tyrosinol, 477
Amphibian oocytes, interspersed poly(A) RNAs are not translatable, 227
Amplification of a 27 nucleotide sequence and the evolution of the albumin : α-fetoprotein ancestral gene, 159
Anderson, D. M. (see also Richter, J. D.), 227
Anderson, N. L. (see also Giometti, C. S.), 109
Antibodies, monoclonal, use in immuno-electron microscopy for the determination of subunit stoichiometry in oligomeric enzymes; there are three α-subunits in the F_1-ATPase of *Escherichia coli*, 131
Arylisothiocyanates, use in group-directed modification of bacteriorhodopsin; labeling, identification of the binding site and topology, 93
Asakura, S. (see also Kato, S.), 463
ATP, interaction with crystalline tyrosyl-tRNA synthetase in the presence of tyrosinol, 477
F_1-ATPase of *Escherichia coli* has three α-subunits; use of monoclonal antibodies in immuno-electron microscopy for the determination of subunit stoichiometry in oligomeric enzymes, 131
Avolio, J., Lebduska, S. & Satir, P., Dynein arm substructure and the orientation of arm–microtubule attachments, 389

Bacteriophage lambda, changes in DNA base sequence due to targeted mutagenesis by ultraviolet light, 273
——, unirradiated, non-targeted mutagenesis in *Escherichia coli* host cells irradiated with ultraviolet light, 293
— T7 RNA polymerase, nucleotide sequence of the gene, 265
Bacteriorhodopsin, group-directed modification by arylisothiocyanates; labeling, identification of the binding site and topology, 93
Beck, E., Jorcano, J. L. & Alonso, A., *Drosophila melanogaster* U1 and U2 small nuclear RNA genes contain common flanking sequences, 539
Bishop, C., Guellaen, G., Geldwerth, D., Fellous, M. & Weissenbach, J., Extensive sequence homologies between Y and other human chromosomes, 403
Blow, D. M. (see also Monteilhet, C.), 477
Bobst, A. M., Kao, S.-C., Toppin, R. C., Ireland, J. C. & Thomas, I. E., Dipsticking the major groove of DNA with enzymatically incorporated spin-labeled deoxyuridines by electron spin resonance spectroscopy, 63
Brick, P. (see also Monteilhet, C.), 477
Bull seminal plasma, secondary structure in the solution conformation of the proteinase inhibitor IIA by n.m.r., 341

Calf satellite DNA I, nature and organization of the sequence variations, 143
Caulobacter crescentus flagellar filament has a right-handed helical form, 125
Chattoraj, D. K. (see also Abeles, A. L.), 293

Chromosomes, human, Y and others, have extensive sequence homologies, 403
Chromosome, yeast 2 μm plasmid DNA, location of DNAase I sensitive cleavage sites. 1
Cleavage sites, DNAase I sensitive, location in the yeast 2 μm plasmid DNA chromosome, 1
— specificity, how it is maintained by signal sequences, 243
Comer, M. M. (see also Wu, T.-H.), 177
Cross-bridge attachment and changes in the lateral filament spacing of skinned muscle fibres, 15
Crystalline tyrosyl-tRNA synthetase, interaction with adenosine, adenosine monophosphate, adenosine triphosphate and pyrophosphate in the presence of tyrosinol, 477
Crystallographic determination of local mobility of nucleic acids; RNA and *B* form DNA, 361

Darlix, J.-L. (see also Méric, C.), 531
Davidson, E. H. (see also Richter, J. D.), 227
Deletion analysis of transposon Tn*9*, 523
DNA, *B* form, crystallographic determination of local mobility, 361
— base sequence changes induced by targeted mutagenesis of lambda phage by ultraviolet light, 273
—, dipsticking the major groove with enzymatically incorporated spin-labeled deoxyuridines by electron spin resonance spectroscopy, 63
— sequences, repetitive, interspersed, non-Alu family, in the mouse genome; organization and expression, 419
— supercoiling, length dependence of the free energy, and torsional rigidity of DNA, 75
— torsional rigidity and length dependence of the free energy of DNA supercoiling, 75
— I, satellite, calf, nature and organization of the sequence variations, 143
DNAase I sensitive cleavage sites, location in the yeast 2 μm plasmid DNA chromosome, 1
DNA-binding proteins, two, and specificity of translational regulation, 211
Deoxyuridines, spin-labeled, enzymatically incorporated, use in dipsticking the major groove of DNA by electron spin resonance spectroscopy, 63
Dimensions, molecular, of the SS and FF phenotypes of human placental alkaline phosphatase; rotary shadowing and negative staining electron microscope measurements, 253
Dipsticking the major groove of DNA with enzymatically incorporated spin-labeled deoxyuridines by electron spin resonance spectroscopy, 63
Drosophila melanogaster U1 and U2 small nuclear RNA genes contain common flanking sequences, 539
Dunn, J. J. (see also Moffatt, B. A.), 265
Dynein arm substructure and the orientation of arm–microtubule attachments, 389

Ehrenfeld, E. (see also Richards, O. C.), 325
Ehrig, K. (see also Lünsdorf, H.), 131
Electron microscope analysis of RNA cross-linked *in vivo* with psoralen derivative; structure of poliovirus replicative intermediate RNA, 325
— microscopy, rotary shadowing and negative staining, and measurements of the molecular dimensions of the SS and FF phenotypes of human placental alkaline phosphatase, 253
— spin resonance spectroscopy, use in dipsticking the major groove of DNA with enzymatically incorporated spin-labeled deoxyuridines, 63
Electrophoretic variant of a human fibroblast protein with characteristics of smooth muscle tropomyosin, 109
Environment of ionizable groups in globular proteins, 515
Enzyme-probe method to detect structural changes in the myosin rod, 35
Enzymes, oligomeric, use of monoclonal antibodies in immuno-electron microscopy for the determination of subunit stoichiometry; there are three α-subunits in the F_1-ATPase of *Escherichia coli*, 131
Escherichia coli F_1-ATPase has three α-subunits; use of monoclonal antibodies in immuno-electron microscopy for the determination of subunit stoichiometry in oligomeric enzymes, 131
— — host cells irradiated with ultraviolet light, and non-targeted mutagenesis of unirradiated lambda phage, 293
— —, multiple IS*10* rearrangements, 437
— —, polymorphic transition of the flagellar polyhook, 463
— —, transcription of a gene cluster coding for two aminoacyl-tRNA synthetases and an initiation factor, 177
Evolution of the albumin: α-fetoprotein ancestral gene from the amplification of a 27 nucleotide sequence, 159
Expression and organization of non-Alu family interspersed repetitive DNA sequences in the mouse genome, 419

Fagrelius, T. J. & Livingston, D. M., Location of DNAase I sensitive cleavage sites in the yeast 2 μm plasmid DNA chromosome, 1
Fellous, M. (see also Bishop, C.), 403
α-Fetoprotein : albumin gene, evolution from the amplification of a 27 nucleotide sequence, 159
— gene, determination of the likelihood of repetitive sequences, 174
Fibroblast protein, human, an electrophoretic variant with characteristics of smooth muscle tropomyosin, 109
Filament lattice of skinned muscle fibres, changes in the lateral spacing when cross-bridges attach, 15
Fishman, W. H. (see also Takeya, M.), 253
Flagellar filament of *Caulobacter crescentus* has a right-handed helical form, 125
— polyhook from *Escherichia coli* and *Salmonella typhimurium*, polymorphic transition, 463
Free energy of DNA supercoiling, length dependence, and torsional rigidity of DNA, 75
Friedl, P. (see also Lünsdorf, H.), 131
Fulford, W. & Model, P., Specificity of translational regulation by two DNA-binding proteins, 211

Geldwerth, (see also Bishop, C.), 403
Gene, ancestral albumin : α-fetoprotein, evolution from the amplification of a 27 nucleotide sequence, 159
— for bacteriophage T7 RNA polymerase, the nucleotide sequence, 265
— cluster coding for two aminoacyl-tRNA synthetases and an initiation factor in *Escherichia coli*, transcription, 177
Genes, U1 and U2 small nuclear RNA, of *Drosophila melanogaster*, contain common flanking sequences, 539
Genome, mouse, organization and expression of non-Alu family interspersed repetitive DNA sequences, 419
Genzel, L. (see also Poglitsch, A.), 137
Giometti, C. S. & Anderson, N. L., An electrophoretic variant of a human fibroblast protein with characteristics of smooth muscle tropomyosin, 109
Globular proteins, the environment of ionizable groups, 515
Goldman, Y. E. (see also Matsubara, I.), 15
Guellaen, G. (see also Bishop, C.), 403

Harrington, W. F. (see also Ueno, H.), 35
Helical form, right-handed, of *Caulobacter crescentus* flagellar filament, 125
Heller, D., Jackson, M. & Leinwand, L., Organization and expression of non-Alu family interspersed repetitive DNA sequences in the mouse genome, 419
Holbrook, S. R. & Kim, S.-H., Local mobility of nucleic acids as determined from crystallographic data. I. RNA and *B* form DNA, 361
Honig, B. (see also Rashin, A. A.), 515
Horowitz, D. S. & Wang, J. C., Torsional rigidity of DNA and length dependence of the free energy of DNA supercoiling, 75
Human chromosomes, Y and others, have extensive sequence homologies, 403
— fibroblast protein, an electrophoretic variant with characteristics of smooth muscle tropomyosin, 109
— placental alkaline phosphatase, molecular dimensions of the SS and FF phenotypes measured by rotary shadowing and negative staining electron microscopy, 253
Hutchinson, F. (see also Wood, R. D.), 273, 293
Hydrated lysozyme, picosecond relaxations observed by mm-wave spectroscopy, 137

Immuno-electron microscopy and the use of monoclonal antibodies for the determination of subunit stoichiometry in oligomeric enzymes; there are three α-subunits in the F_1-ATPase of *Escherichia coli*, 131
Initiation factor and two aminoacyl-tRNA synthetases in *Escherichia coli*, transcription of the coding gene cluster, 177
Interspersed poly(A) RNAs of amphibian oocytes are not translatable, 227
— repetitive DNA sequences, non-Alu family, in the mouse genome; organization and expression, 419
Ionizable groups in globular proteins; their environment, 515
Ireland, J. C. (see also Bobst, A. M.), 63
IS*10* multiple rearrangements in *Escherichia coli*, 437

Jackson, M. (see also Heller, D.), 419

Jense, H. G. (see also Richards, O. C.), 325
Jorcano, J. L. (see also Beck, E.), 539

Kao, S.-C. (see also Bobst, A. M.), 63
Kato, S., Okamoto, M. & Asakura, S., Polymorphic transition of the flagellar polyhook from *Escherichia coli* and *Salmonella typhimurium*, 463
Kim, S.-H. (see also Holbrook, S. R.), 361
Kleckner, N. (see also Raleigh, E. A.), 437
Klier, F. G. (see also Takeya, M.), 253
Koyasu, S. & Shirakihara, Y., *Caulobacter crescentus* flagellar filament has a right-handed helical form, 125
Kremer, F. (see also Poglitsch, A.), 137

λ (see also Bacteriophage), 273, 293
Lebduska, S. (see also Avolio, J.), 389
Leinwand, L. (see also Heller, D.), 419
Litwin, S. (see also Young, P. R.), 174
Livingston, D. M. (see also Fagrelius, T. J.), 1
Local mobility of nucleic acids as determined from crystallographic data; RNA and *B* form DNA, 361
Lünsdorf, H., Ehrig, K., Friedl, P. & Schairer, H. U., Use of monoclonal antibodies in immuno-electron microscopy for the determination of subunit stoichiometry in oligomeric enzymes. There are three α-subunits in the F_1-ATPase of *Escherichia coli*, 131
Lysozyme, hydrated, picosecond relaxations observed by mm-wave spectroscopy, 137

Marion, D. (see also Williamson, M. P.), 341
Martin, S. C. (see also Richards, O. C.), 325
Matsubara, I., Goldman, Y. E. & Simmons, R. M., Changes in the lateral filament spacing of skinned muscle fibres when crossbridges attach, 15
Méric, C., Darlix, J.-L. & Spahr, P. F., It is Rous Sarcoma virus protein P12 and not P19 that binds tightly to Rous Sarcoma virus RNA, 531
Microtubule attachments to dynein arm, study of orientation, 389
mm-wave spectroscopic observations of picosecond relaxations in hydrated lysozyme, 137
Mobility, local, of nucleic acids as determined from crystallographic data; RNA and *B* form DNA, 361
Model, P. (see also Fulford, W.), 211
Modification, group-directed, of bacteriorhodopsin by arylisothiocyanates; labeling, identification of the binding site and topology, 93
Moffatt, B. A., Dunn, J. J. & Studier, F. W., Nucleotide sequence of the gene for bacteriophage T7 RNA polymerase, 265
Monoclonal antibodies, use in immuno-electron microscopy for the determination of subunit stoichiometry in oligomeric enzymes; there are three α-subunits in the F_1-ATPase of *Escherichia coli*, 131
Monteilhet, C., Blow, D. M. & Brick, P. Interaction of crystalline tyrosyl-tRNA synthetase with adenosine, adenosine monophosphate, adenosine triphosphate and pyrophosphate in the presence of tyrosinol, 477
Mouse genome, organization and expression of non-Alu family interspersed repetitive DNA sequences, 419
Muscle fibres, skinned, changes in the lateral filament spacing when cross-bridges attach, 15
— tropomyosin characteristics of an electrophoretic variant of a human fibroblast protein, 109
Mutagenesis, non-targeted, of unirradiated lambda phage in *Escherichia coli* host cells irradiated with ultraviolet light, 293
—, targeted, of lambda phage by ultraviolet light, and induced changes in DNA base sequence, 273
Myosin rod, an enzyme-probe method to detect structural changes, 35

Negative staining and rotary shadowing electron microscope measurements of molecular dimensions of the SS and FF phenotypes of human placental alkaline phosphatase, 253
Nuclear magnetic resonance study of secondary structure in the solution conformation of the proteinase inhibitor IIA from bull seminal plasma, 341
Nucleic acid local mobility as determined from crystallographic data; RNA and *B* form DNA, 361
Nucleotide sequence of the gene for bacteriophage T7 RNA polymerase, 265

Okamoto, M. (see also Kato, S.), 463

Oocytes, amphibian, interspersed poly(A) RNAs are not translatable, 227
Organization and expression of non-Alu family interspersed repetitive DNA sequences in the mouse genome, 419

P1 plasmid replication: replicon structure, 307
Pagès, M. J. M. & Roizès, G. P., Nature and organization of the sequence variations in the long-range periodicity calf satellite DNA I, 143
Phenotypes SS and FF of human placental alkaline phosphatase, molecular dimensions measured by rotary shadowing and negative staining electron microscopy, 253
Picosecond relaxations in hydrated lysozyme observed by mm-wave spectroscopy, 137
Placental alkaline phosphatase, human, molecular dimensions of the SS and FF phenotypes measured by rotary shadowing and negative staining electron microscopy, 253
Plasmid DNA chromosome, 2 μm, yeast, location of DNAase I sensitive cleavage sites, 1
— P1 replication; replicon structure, 307
Poglitsch, A., Kremer, F., Genzel, L., Picosecond relaxations in hydrated lysozyme observed by mm-wave spectroscopy, 137
Poliovirus replicative intermediate RNA, structure; electron microscope analysis of RNA cross-linked *in vivo* with psoralen derivative, 325
Poly(A) RNAs, interspersed, of amphibian oocytes are not translatable, 227
Polyhook, flagellar, from *Escherichia coli* and *Salmonella typhimurium*; polymorphic transition, 463
Protein, fibroblast, human, an electrophoretic variant with characteristics of smooth muscle tropomyosin, 109
— P12 of Rous Sarcoma virus, and not P19, binds tightly to Rous Sarcoma virus RNA, 531
Proteins, DNA-binding, two, and specificity of translational regulation, 211
—, globular, the environment of ionizable groups, 515
—, recognition of super-secondary structure, 487
Proteinase inhibitor IIA from bull seminal plasma, secondary structure in the solution conformation by n.m.r., 341
Psoralen derivative cross-linked *in vivo* with RNA, electron microscope analysis; structure of poliovirus replicative intermediate RNA, 325
Pyrophosphate, interaction with crystalline tyrosyl-tRNA synthetase in the presence of tyrosinol, 477

Raleigh, E. A. & Kleckner, N., Multiple IS*10* rearrangements in *Escherichia coli*, 437
Rashin, A. A. & Honig, B., On the environment of ionizable groups in globular proteins, 515
Rearrangements, IS*10*, multiple, in *Escherichia coli*, 437
Regulation, translational, by two DNA-binding proteins; specificity, 211
Relaxations, picosecond, in hydrated lysozyme observed by mm-wave spectroscopy, 137
Repetitive DNA sequences, interspersed, non-Alu family, in the mouse genome; organization and expression, 419
— sequences, determination of likelihood in the α-fetoprotein gene, 174
Replication, P1 plasmid; replicon structure, 307
Replicative intermediate RNA of poliovirus, structure; electron microscope analysis of RNA cross-linked *in vivo* with psoralen derivative, 325
RNA, crystallographic determination of local mobility, 361
— genes U1 and U2 of *Drosophila melanogaster* contain common flanking sequences, 539
—, intermediate, replicative, poliovirus, a study of structure; electron microscope analysis of RNA cross-linked *in vivo* with psoralen derivative, 325
— polymerase, T7, nucleotide sequence of the gene, 265
—, Rous Sarcoma virus, binds tightly to Rous Sarcoma virus protein P12 and not P19, 531
RNAs, poly(A), interspersed, of amphibian oocytes are not translatable, 227
Richards, O. C., Martin, S. C., Jense, H. G. & Ehrenfeld, E., Structure of poliovirus replicative intermediate RNA. Electron microscope analysis of RNA cross-linked *in vivo* with psoralen derivative, 325
Richter, J. D., Smith, L. D., Anderson, D. M. & Davidson, E. H., Interspersed poly(A) RNAs of amphibian oocytes are not translatable, 227
Rickli, E. E. (see also Sigrist, H.), 93
Rigidity, torsional, of DNA and length dependence of the free energy of DNA supercoiling, 75
Roizès, G. P. (see also Pagès, M. J. M.), 143
Rotary shadowing and negative staining electron microscope measurements of molecular dimensions of the SS and FF phenotypes of human placental alkaline phosphatase, 253
Rous Sarcoma virus (see also Virus), 531

Salmonella typhimurium, polymorphic transition of the flagellar polyhook, 463
Satellite DNA I, calf, nature and organization of the sequence variations, 143
Satir, P. (see also Avolio, J.), 389
Schairer, H. U. (see also Lünsdorf, H.), 131
Schaller, J. (see also Sigrist, H.), 93
Secondary structure, an algorithm to compare predictions, 512
Seminal plasma, bull, secondary structure in the solution conformation of the proteinase inhibitor IIA by n.m.r., 341
Sequence amplification and the evolution of the albumin : α-fetoprotein ancestral gene, 159
— homologies, extensive, between Y and other human chromosomes, 403
—, nucleotide, of the gene for bacteriophage T7 RNA polymerase, 265
— variations in the long-range periodicity calf satellite DNA I, nature and organization, 143
Sequences, DNA, repetitive, interspersed, non-Alu family, in the mouse genome; organization and expression, 419
—, flanking, common, of *Drosophila melanogaster* U1 and U2 small nuclear RNA genes, 539
—, signal, how they maintain cleavage specificity, 243
Shirakihara, Y. (see also Koyasu, S.), 125
Signal sequences, how they maintain cleavage specificity, 243
Sigrist, H., Allegrini, P. R., Stauffer, K., Schaller, J., Abdulaev, N. G., Rickli, E. E. & Zahler, P., Group-directed modification of bacteriorhodopsin by arylisothiocyanates. Labeling, identification of the binding site and topology, 93
Simmons, R. M. (see also Matsubara, I.), 15
Skinned muscle fibres, changes in the lateral filament spacing when cross-bridges attach, 15
Skopek, T. R. (see also Wood, R. D.), 273
Small nuclear RNA genes U1 and U2 of *Drosophila melanogaster* contain common flanking sequences, 539
Smith, L. D. (see also Richter, J. D.), 227
Snyder, K. M. (see also Abeles, A. L.), 293
Spahr, P.-F. (see also Méric, C.), 531
Specificity, cleavage, how it is maintained by signal sequences, 243
— of translational regulation by two DNA-binding proteins, 211
Stauffer, K. (see also Sigrist, H.), 93
Stein, P. L. (see also Wu, T.-H.), 177
Stoichiometry, subunit, in oligomeric enzymes, determination using monoclonal antibodies and immuno-electron microscopy; there are three α-subunits in the F_1-ATPase of *Escherichia coli*, 131
Structural changes in the myosin rod, an enzyme-probe method of detection, 35
Structure of the dynein arm and the orientation of arm-microtubule attachments, 389
poliovirus replicative intermediate RNA. Electron microscope analysis of RNA cross-linked *in vivo* with psoralen derivative, 325
— — replicon in P1 plasmid replication, 307
—, secondary, an algorithm to compare predictions, 512
—, —, in the solution conformation of the proteinase inhibitor IIA from bull seminal plasma by n.m.r., 341
—, super-secondary, recognition in proteins, 487
Studier, F. W. (see also Moffatt, B. A.), 265
Subunit stoichiometry in oligomeric enzymes, determination using monoclonal antibodies and immuno-electron microscopy; there are three α-subunits in the F_1-ATPase of *Escherichia coli*, 131
Supercoiling, DNA, length dependence of the free energy, and the torsional rigidity of DNA, 75
Super-secondary structure recognition in proteins, 487

T7 (see also Bacteriophage), 265
Takeya, M., Klier, F. G. & Fishman, W. H., Molecular dimensions of the SS and FF phenotypes of human placental alkaline phosphatase. Rotary shadowing and negative staining electron microscope measurements, 253
Targeted mutagenesis of lambda phage by ultraviolet light, and induced changes in DNA base sequence, 273
Taylor, W. R., An algorithm to compare secondary structure predictions (*Appendix* to Taylor, W. R. & Thornton, J. M., 487), 512
— & Thornton, J. M., Recognition of super-secondary structure in proteins, 487
Thomas, I. E. (see also Bobst, A. M.), 63
Thornton, J. M. (see also Taylor, W. R.), 487

LGHMAN, S. M. (see also ALEXANDER, F.), 159
9, a deletion analysis, 523
PPIN, R. C. (see also BOBST, A. M.), 63
rsional rigidity of DNA and length dependence of the free energy of DNA supercoiling, 75
anscription of a gene cluster coding for two aminoacyl-tRNA synthetases and an initiation factor in *Escherichia coli*, 177
ansition, polymorphic, of the flagellar polyhook from *Escherichia coli* and *Salmonella typhimurium*, 463
anslation of interspersed poly(A) RNAs of amphibian oocytes is not possible, 227
anslational regulation by two DNA-binding proteins, specificity, 211
ansposon Tn*9*, a deletion analysis, 523
opomyosin, smooth muscle, characteristics of an electrophoretic variant of a human fibroblast protein, 109
rosinol, presence in the interaction of crystalline tyrosyl-tRNA synthetase with adenosine, adenosine monophosphate, adenosine triphosphate and pyrophosphate, 477

and U2 *Drosophila melanogaster* small nuclear RNA genes contain common flanking sequences, 539
ENO, H. & HARRINGTON, W. F., An enzyme-probe method to detect structural changes in the myosin rod, 35
traviolet light irradiation of *Escherichia coli* host cells, and non-targeted mutagenesis of unirradiated lambda phage, 293
—, use in inducing changes in DNA base sequence by targeted mutagenesis of lambda phage, 273

rus, Rous Sarcoma: it is protein P12 and not P19 that binds tightly to Rous Sarcoma virus RNA, 531
N HEIJNE, G., How signal sequences maintain cleavage specificity, 243

ANG, J. C. (see also HOROWITZ, D. S.), 75
EISSENBACH, J. (see also BISHOP, C.), 403
ILLIAMSON, M. P., MARION, D. & WÜTHRICH, K., Secondary structure in the solution conformation of the proteinase inhibitor IIA from bull seminal plasma by nuclear magnetic resonance, 341
OOD, D. L. (see also WU, T.-H.), 177
OOD, R. D. & HUTCHINSON, F., Non-targeted mutagenesis of unirradiated lambda phage in *Escherichia coli* host cells irradiated with ultraviolet light, 293
, SKOPEK, T. R. & HUTCHINSON, F., Changes in DNA base sequence induced by targeted mutagenesis of lambda phage by ultraviolet light, 273
U, T.-H., WOOD, D. L., STEIN, P. L. & COMER, M. M., Transcription of a gene cluster coding for two aminoacyl-tRNA synthetases and an initiation factor in *Escherichia coli*, 177
ÜTHRICH, K. (see also WILLIAMSON, M. P.), 341

and other human chromosomes have extensive sequence homologies, 403
east 2 μm plasmid DNA chromosome, location of DNAase I sensitive cleavage sites, 1
OUNG, P. R. & LITWIN, S., Determination of the likelihood of repetitive sequences in the α-fetoprotein gene (*Appendix* to ALEXANDER, F., *et al.*, 159), 174
(see also ALEXANDER, F.), 159

AHLER, P. (see also SIGRIST, H.), 93